

## ColoredTaq DNA Polymerase

Cat. No.	Pack size	Conc.	Supplied buffer
E021	500 u	2.5 u/μl	x10 Taq Buffer, 25mM Mg <sup>2+</sup> x10 Taq Buffer, without Mg <sup>2+</sup> 25 mM MgCl <sub>2</sub>
E022	2x2'500 u	2.5 u/μl	x10 Taq Buffer, 25mM Mg <sup>2+</sup> x10 Taq Buffer, without Mg <sup>2+</sup> 25 mM MgCl <sub>2</sub>



### Description:

ColoredTaq polymerase is a variant of Taq polymerase. There is a special neutral dye added to the enzyme to make it possible to track the movement of samples in a gel. The reaction mixture with ColoredTaq polymerase is "ready-to-load" on the gel (no additional dye loading is required).

Taq polymerase is a thermostable enzyme with mass of approx. 94kDa. The source of the enzyme is a strain of *E.coli* with plasmid encoding the sequence of Taq DNA polymerase. The enzyme has activity maximum at 70-74 °C (although the enzyme is able to work at lower temperatures of about 25 °C). Enzyme's half-life at 94 °C is at least 45 min. The enzyme catalyzes DNA synthesis in 5'→3' direction in the presence of Mg<sup>2+</sup> and at the appropriate pH. It also has a 5'→3' exonuclease activity (this activity is realized only on double-stranded DNA). The presence of the 5'→3' exonuclease activity makes the enzyme suitable for use in RealTime PCR. Taq polymerase adds extra adenines at the 3'-end of PCR products. It does not create interferences with cloning. If it is necessary to obtain PCR products without extra adenines we recommend using Pfu polymerase. The maximum length of PCR products that can be obtained using Taq polymerase is approx. 5,000 bp.

### Applications:

- EndPoint PCR

### Buffers:

- **x10 Taq Buffer, 25 mM Mg<sup>2+</sup>**  
700 mM Tris-HCl, pH 8.8 at 25 °C,  
166 mM (NH<sub>4</sub>)<sub>2</sub>SO<sub>4</sub>,  
25 mM MgCl<sub>2</sub>
- **x10 Taq Buffer, without Mg<sup>2+</sup>**  
700 mM Tris-HCl, pH 8.8 at 25 °C,  
166 mM (NH<sub>4</sub>)<sub>2</sub>SO<sub>4</sub>

### Storage buffer:

50% glycerol (v/v),  
20 mM Tris-HCl, pH 7.5 at 25 °C,  
100 mM NaCl,  
0.1 mM EDTA,  
2 mM DTT,  
0.2% Tween 20

### Concentration:

2.5 u/μl

### Unit definition:

One unit is defined as the amount of enzyme required to catalyze the incorporation of 10 nmol of dNTPs into an acid-insoluble form in 30 minutes at 70 °C.

### Quality control:

The enzyme is tested for the absence of nicking activities, of endo-, exodeoxyribonucleases and ribonucleases. Activity and thermostability tested in PCR.

### Storage conditions:

Store at -20 °C

### Shipping conditions:

No special conditions required.

### Safety and precautions:

Avoid contact with eyes, skin and clothing.  
In case of contact with eyes, rinse immediately with plenty of water for at least 15 minutes and seek medical advice. Make the rinsing more thorough by separating the eyelids with fingers. If symptoms such as redness and irritation persist, seek medical attention.

In case of contact with skin, immediately wash the skin with soap and plenty amounts of water.

**Working with ColoredTaq DNA Polymerase:**



1. Test tube with all components added except the polymerase




2. Adding ColoredTaq Polymerase. It is more dense than water




3. ColoredTaq does not mix with the solution and goes to the bottom



4. After shaking on vortex ColoredTaq mixes with the tube contents and is ready to work



5. When loading the reaction mix on gel it also goes down making the procedure more simple



6. No additional dye needed because ColoredTaq allows easy visualisation of the product migration in gel

**Recommended PCR protocol:**

*(the protocol can be modified according to use of another components or volume)*

Mix the following components (total volume - 25 µl):	
x10 Taq Buffer, 25mM Mg <sup>2+</sup> or x10 Taq Buffer, without Mg <sup>2+</sup> 25 mM MgCl <sub>2</sub>	2.5 µl
2.5 mM dNTP Mix	2.5 µl
Primer 1 (10 pmol/µl)	0.5 µl
Primer 2 (10 pmol/µl)	0.5 µl
Template DNA (10 ng)	2.0 µl
ColoredTaq polymerase (2.5 u/µl)	0.5 - 1.0 µl
ddH <sub>2</sub> O	up to 25 µl

PCR cycles:

Step	Temp.	Time
Initial denaturation	94 °C	1 min
Denaturation	94 °C	10 sec
Annealing	60 °C	10 sec
Elongation	72 °C	10 sec

30 cycles

**Note:** Annealing temperature, step duration and cycles number can be modified according to the actual tasks.